

# Math 5510 - Partial Differential Equations

## Fourier Transforms for PDEs - Part B

**Ahmed Kaffel**

`<ahmed.kaffel@marquette.edu>`

Department of Mathematical and Statistical Sciences  
Marquette University

<https://www.mscsnet.mu.edu/~ahmed/teaching.html>

Spring 2021

# Outline

- 1 Heat Equation and Fourier Transforms
  - Fundamental Solution and  $\delta(x)$
  - Example
  
- 2 Fourier Transforms of Derivatives
  - Heat Equation
  - Convolution

# Heat Equation and Fourier Transforms

We showed that  $e^{-i\omega x}e^{-k\omega^2 t}$  solve the *heat equation*,  $u_t = ku_{xx}$ , so

$$u(x, t) = \int_{-\infty}^{\infty} c(\omega)e^{-i\omega x}e^{-k\omega^2 t} d\omega.$$

The **IC** is satisfied if:

$$f(x) = \int_{-\infty}^{\infty} c(\omega)e^{-i\omega x} d\omega.$$

From the definition of the *Fourier transform*, the above equation is a *Fourier integral* representation of  $f(x)$  with

$$c(\omega) = \frac{1}{2\pi} \int_{-\infty}^{\infty} f(x)e^{i\omega x} dx.$$

# Heat Equation and Fourier Transforms

The *Fourier coefficient* can be inserted into the solution:

$$u(x, t) = \int_{-\infty}^{\infty} \left[ \frac{1}{2\pi} \int_{-\infty}^{\infty} f(s) e^{i\omega s} ds \right] e^{-i\omega x} e^{-k\omega^2 t} d\omega.$$

Interchanging the order of integration gives:

$$u(x, t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} f(s) \left[ \int_{-\infty}^{\infty} e^{-k\omega^2 t} e^{-i\omega(x-s)} d\omega \right] ds.$$

However, the *inverse Fourier transform* of  $e^{-k\omega^2 t}$

$$g(x) = \int_{-\infty}^{\infty} e^{-k\omega^2 t} e^{-i\omega x} d\omega = \sqrt{\frac{\pi}{kt}} e^{-x^2/4kt}.$$

# Heat Equation and Fourier Transforms

We insert the information above into the solution and obtain:

$$u(x, t) = \int_{-\infty}^{\infty} f(s) \left[ \frac{1}{\sqrt{4\pi kt}} e^{-(x-s)^2/4kt} \right] ds.$$

It follows that each initial temperature “*influences*” the temperature at time  $t$  according to the *Influence function*, which is related to the *Green's functions* last section:

$$G(x, t; s, 0) = \frac{1}{\sqrt{4\pi kt}} e^{-(x-s)^2/4kt}.$$

This *Influence function* has problems near  $t = 0$ .

Dirac Delta function,  $\delta(x)$ 

Define the function:

$$f(x, a) = \begin{cases} 0, & |x| > a, \\ \frac{1}{2a}, & |x| < a. \end{cases}$$

The *Dirac delta function* satisfies:

$$\lim_{a \rightarrow 0} f(x, a) = \delta(x).$$


With regards to our *Heat problem*, we see that as  $t \rightarrow 0$  the *influence* is concentrated locally:

$$\lim_{t \rightarrow 0^+} \frac{1}{\sqrt{4\pi kt}} e^{-(x-s)^2/4kt} = \delta(x-s).$$

# Fundamental Solution

**Fundamental Solution:** Suppose all the heat is concentrated at the origin,  $u(x, 0) = \delta(x)$ , then

$$u(x, t) = \int_{-\infty}^{\infty} \delta(s) \left[ \frac{1}{\sqrt{4\pi kt}} e^{-(x-s)^2/4kt} \right] ds = \frac{1}{\sqrt{4\pi kt}} e^{-x^2/4kt}.$$



# Heat Equation and Fourier Transforms

**Example:** Consider the infinite rod satisfying the *heat equation*:

$$\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2}, \quad t > 0, \quad -\infty < x < \infty,$$

with **IC**

$$u(x, 0) = f(x) = \begin{cases} 0, & x < 0, \\ 100, & x > 0. \end{cases}$$

From above the solution satisfies:

$$\begin{aligned} u(x, t) &= \int_{-\infty}^{\infty} f(s) \left[ \frac{1}{\sqrt{4\pi kt}} e^{-(x-s)^2/4kt} \right] ds, \\ &= \frac{100}{\sqrt{4\pi kt}} \int_0^{\infty} e^{-(x-s)^2/4kt} ds. \end{aligned}$$



# Heat Equation and Fourier Transforms

With the change of dummy variables in the integral,  
 $z = (s - x)/\sqrt{4kt}$ , the solution can be written:

$$\begin{aligned}u(x, t) &= \frac{100}{\sqrt{4\pi kt}} \int_0^\infty e^{-(x-s)^2/4kt} ds, \\&= \frac{100}{\sqrt{\pi}} \int_{-x/\sqrt{4kt}}^\infty e^{-z^2} dz, \\&= \frac{100}{\sqrt{\pi}} \left[ \int_0^\infty e^{-z^2} dz + \int_0^{x/\sqrt{4kt}} e^{-z^2} dz \right],\end{aligned}$$

by the evenness of  $e^{-z^2}$ .



Thus, we can write the solution:

$$\begin{aligned}u(x, t) &= 50 + \frac{100}{\sqrt{\pi}} \int_0^{x/\sqrt{4kt}} e^{-z^2} dz, \\&= 50 \left( 1 + \operatorname{erf} \left( \frac{x}{\sqrt{4kt}} \right) \right).\end{aligned}$$

# Heat Equation and Fourier Transforms

The *temperature* spreads by *diffusion*.

The thermal energy spreads with infinite propagation speed.



# Heat Equation and Fourier Transforms

Below is the MatLab code for the previous figures for **Heat Propagation**.

```
1 % Solutions to the heat flow equation
2 % on one-dimensional rod
3 % Fourier Transform solution
4 format compact;
5 tfin = 100;           % final time
6 xwid = 10;
7 k = 1;               % heat capacitance
8 NptsT=151;           % number of t pts
9 NptsX=151;           % number of x pts
10 t=linspace(0,tfin,NptsT);
11 x=linspace(-xwid,xwid,NptsX);
12 [T,X]=meshgrid(t,x);
13
14 figure(1)
```

## Heat Equation and Fourier Transforms

```
15 clf
16 U = 50*(1 + erf(X./(sqrt(4*k*T)))); % Temperature(n)
17
18 set(gca,'FontSize',[12]);
19 surf(T,X,U);
20 shading interp
21 colormap(jet)
22 xlabel('$t$', 'FontSize',12, 'interpreter','latex');
23 ylabel('$x$', 'FontSize',12, 'interpreter','latex');
24 zlabel('$u(x,t)$', 'FontSize',12, 'interpreter','latex');
25 axis tight
26 view([30 12])
27 print -dpng heatFT1.png
28 print -depsc heatFT1.eps
```

## Heat Equation and Fourier Transforms

```
30 figure(2)
31 clf
32
33 set(gca, 'FontSize', [12]);
34 surf(T, X, U);
35 shading interp
36 colormap(jet)
37 view([0 90])           %create 2D color map of ...
                        temperature
38 xlabel('$t$', 'Fontsize', 12, 'interpreter', 'latex');
39 ylabel('$x$', 'Fontsize', 12, 'interpreter', 'latex');
40 zlabel('$u(x,t)$', 'Fontsize', 12, 'interpreter', 'latex');
41 axis tight
42 colorbar
43 set(gca, 'FontSize', [12]);
44 print -dpng heatFT2.png
45 print -depsc heatFT2.eps
```

# Fourier Transforms of Derivatives

Again consider the *Heat equation*:

$$\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2}, \quad t > 0, \quad -\infty < x < \infty,$$

with **IC**,  $u(x, 0) = f(x)$ .

*Separation of variables* motivated the *Fourier transform*.

Now solve this directly with *Fourier transform*.

Define

$$\mathcal{F}[u] = \bar{U}(\omega, t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} u(x, t) e^{i\omega x} dx$$

be the *Fourier transform* of  $u(x, t)$ .

# Fourier Transforms of Derivatives

Take the partial with respect to  $t$ ,

$$\mathcal{F}\left[\frac{\partial u}{\partial t}\right] = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\partial u}{\partial t} e^{i\omega x} dx = \frac{\partial}{\partial t} \left[ \frac{1}{2\pi} \int_{-\infty}^{\infty} u(x, t) e^{i\omega x} dx \right] = \frac{\partial}{\partial t} \bar{U}(\omega, t).$$

The *spatial Fourier transform* of a time derivative equals the time derivative of the *Fourier transform*.

Now consider the partial with respect to  $x$

$$\mathcal{F}\left[\frac{\partial u}{\partial x}\right] = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\partial u}{\partial x} e^{i\omega x} dx = \frac{ue^{i\omega x}}{2\pi} \Big|_{-\infty}^{\infty} - \frac{i\omega}{2\pi} \int_{-\infty}^{\infty} u(x, t) e^{i\omega x} dx.$$

If  $\lim_{x \rightarrow \pm\infty} u(x, t) = 0$ , then the endpoints vanish and

$$\mathcal{F}\left[\frac{\partial u}{\partial x}\right] = -i\omega \mathcal{F}[u] = -i\omega \bar{U}(\omega, t).$$

# Fourier Transforms of Derivatives

Similarly, *Fourier transforms* of higher derivatives may be obtained:

$$\mathcal{F} \left[ \frac{\partial^2 u}{\partial x^2} \right] = -i\omega \mathcal{F} \left[ \frac{\partial u}{\partial x} \right] = (-i\omega)^2 \bar{U}(\omega, t) = -\omega^2 \bar{U}(\omega, t).$$

In general, the *Fourier transform* of the  $n^{\text{th}}$  derivative of a function with respect to  $x$  equals  $(-i\omega)^n$  time the *Fourier transform* of the function, assuming that  $u(x, t) \rightarrow 0$  sufficiently fast as  $x \rightarrow \pm\infty$ .

From the properties of the *Fourier transforms* of the derivatives, the *Fourier transform* of the *heat equation* becomes:

$$\frac{\partial}{\partial t} \bar{U}(\omega, t) = -k\omega^2 \bar{U}(\omega, t).$$



# Fourier Transforms of Derivatives

The *Fourier transform* acting on the temperature function,  $u(x, t)$ , converts the linear partial differential equation with constant coefficients into an ordinary differential equation, since the spatial derivatives are transformed into algebraic multiples of the transform.

Since

$$\frac{\partial}{\partial t} \bar{U}(\omega, t) = -k\omega^2 \bar{U}(\omega, t),$$

the solution becomes

$$\bar{U}(\omega, t) = c(\omega)e^{-k\omega^2 t},$$

where the arbitrary constant may depend on  $\omega$ .

The function  $c(\omega)$  comes from the **IC**,  $f(x)$ , so

$$c(\omega) = \frac{1}{2\pi} \int_{-\infty}^{\infty} f(x)e^{i\omega x} dx,$$

which gives the same result as obtained by *separation of variables*.

# Convolution

The solution of the *heat equation* is the product of two functions of  $\omega$ ,

$$\bar{U}(\omega, t) = c(\omega)e^{-k\omega^2 t}.$$

Suppose  $F(\omega)$  and  $G(\omega)$  are *Fourier transforms* of  $f(x)$  and  $g(x)$ :

$$F(\omega) = \frac{1}{2\pi} \int_{-\infty}^{\infty} f(x)e^{i\omega x} dx \quad G(\omega) = \frac{1}{2\pi} \int_{-\infty}^{\infty} g(x)e^{i\omega x} dx$$

$$f(x) = \int_{-\infty}^{\infty} F(\omega)e^{-i\omega x} d\omega \quad g(x) = \int_{-\infty}^{\infty} G(\omega)e^{-i\omega x} d\omega$$

We need to find  $h(x)$  where the *Fourier transform* of  $H(\omega)$  satisfies

$$H(\omega) = F(\omega)G(\omega).$$

# Convolution

Note that

$$\begin{aligned}h(x) &= \int_{-\infty}^{\infty} H(\omega)e^{-i\omega x} d\omega = \int_{-\infty}^{\infty} F(\omega)G(\omega)e^{-i\omega x} d\omega, \\&= \frac{1}{2\pi} \int_{-\infty}^{\infty} F(\omega) \left[ \int_{-\infty}^{\infty} g(s)e^{i\omega s} ds \right] e^{-i\omega x} d\omega, \\&= \frac{1}{2\pi} \int_{-\infty}^{\infty} g(s) \left[ \int_{-\infty}^{\infty} F(\omega)e^{-i\omega(x-s)} d\omega \right] ds, \\h(x) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} g(s)f(x-s) ds, \\&= \frac{1}{2\pi} \int_{-\infty}^{\infty} f(w)g(x-w) dw.\end{aligned}$$

This is the **convolution** of  $f(x)$  and  $g(x)$  usually denoted

$$g * f = f * g$$

# Convolution and Heat Equation

For the *heat equation*, consider the transform  $\bar{U}(\omega, t)$  of the solution  $u(x, t)$ , where

$$\bar{U}(\omega, t) = c(\omega)e^{-k\omega^2 t}.$$

- $c(\omega)$  is the transform of the initial temperature,  $f(x)$ .
- $e^{-k\omega^2 t}$  is the transform of the *fundamental solution*,

$$\sqrt{\frac{\pi}{kt}}e^{-x^2/4kt}.$$

- The **Convolution theorem** gives the solution:

$$u(x, t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} f(s) \sqrt{\frac{\pi}{kt}} e^{-(x-s)^2/4kt} ds.$$

# Convolution and Heat Equation


Enter the **Maple** commands for the graph of  $u(x, t)$

```
u := (x,t) -> (1/(2*Pi))*(int(sqrt(Pi/t)*exp(-(1/4)*(x-s)^2/t), s = -2 .. 2));  
plot3d(u(x,t), x = -10..10, t = 0.0001..20);
```

The **IC** is

$$f(x) = \begin{cases} 1, & |x| < 2, \\ 0, & |x| > 2. \end{cases}$$

This graph shows the *diffusion* of the heat with time.




# Convolution and Heat Equation

This problem can be done in **MatLab** using its integral function, which uses an adaptive quadrature to solve the problem.

```
1 % Solution Heat Equation with FT
2 % Arbitrary f(x)
3
4 N1 = 201; N2 = 201;
5 tv = linspace(0.0001,20,N1);
6 xv = linspace(-10,10,N2);
7 [t1,x1] = ndgrid(tv,xv);
8 f = @(s,c) sqrt(pi/c(1))*exp(-(c(2)-s).^2/(4*c(1)));
9
10 for i = 1:N1
11     for j = 1:N2
12         c = [t1(i,j),x1(i,j)];
13         U(i,j) = ...
14             (1/(2*pi))*integral(@(s)f(s,c),-2,2);
15     end
16 end
```

## Convolution and Heat Equation

```
17 set(gca,'FontSize',[12]);
18 surf(t1,x1,U);
19 shading interp
20 colormap(jet)
21 xlabel('$t$', 'Fontsize',12, 'interpreter','latex');
22 ylabel('$x$', 'Fontsize',12, 'interpreter','latex');
23 zlabel('$u(x,t)$', 'Fontsize',12, 'interpreter','latex');
24 axis tight
25 view([30 12])
```



# Fourier Transforms for PDEs

The *Fourier Transform* technique for solving PDEs is as follows:

- 1 *Fourier Transform* the PDE in one of the variables, often  $x$ .
- 2 Solve the ODE in the other variable, often  $t$ .
- 3 Apply the ICs, determining the initial *Fourier Transform*.
- 4 Use the **convolution theorem** to obtain the solution.

If the IC is only defined on a finite interval, then often **Maple** can manage the integral and produce a 3D plot.



# Parseval's Identity

Since  $h(x)$  is the inverse of the *Fourier Transform* of  $F(\omega)G(\omega)$ :

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} g(s)f(x-s) ds = \int_{-\infty}^{\infty} G(\omega)F(\omega)e^{-\omega x} d\omega.$$

Since this holds for all  $x$ , it holds for  $x = 0$ , so

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} g(s)f(-s) ds = \int_{-\infty}^{\infty} G(\omega)F(\omega) d\omega.$$

Take  $g^*(x) = f(-x)$  to be the complex conjugate, then

$$\begin{aligned} F(\omega) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} f(x)e^{i\omega x} dx = \frac{1}{2\pi} \int_{-\infty}^{\infty} f(-s)e^{-i\omega s} ds \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} g^*(x)e^{-i\omega x} dx = G^*(\omega). \end{aligned}$$

# Parseval's Identity

**Parseval's Identity:**

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} g(x)g^*(x) dx = \int_{-\infty}^{\infty} G(\omega)G^*(\omega) d\omega,$$

or equivalently,

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} |g(x)|^2 dx = \int_{-\infty}^{\infty} |G(\omega)|^2 d\omega.$$

Energy is often proportional to  $|g(x)|^2$ , so

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} |g(x)|^2 dx$$

is the *total energy*.

The quantity  $|G(\omega)|^2$  represents the energy per unit wave number, which is the *spectral energy density*.

The *Fourier Transform*,  $G(\omega)$ , of a function  $g(x)$  is a complex quantity whose magnitude squared is the *spectral energy density* (or amount of energy per unit wave number).