Math 3100

#### Section 1.4: The Matrix Equation Ax = b.

**Definition** Let *A* be an  $m \times n$  matrix whose columns are the vectors  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$  (each in  $\mathbb{R}^m$ ), and let **x** be a vector in  $\mathbb{R}^n$ . Then the product of *A* and **x**, denoted by

#### Ax

is the linear combination of the columns of A whose weights are the corresponding entries in **x**. That is

$$A\mathbf{x} = x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_n\mathbf{a}_n.$$

(Note that the result is a vector in  $\mathbb{R}^{m!}$ )

イロト 不得 トイヨト イヨト

Find the product Ax. Simplify to the extent possible.

The product 
$$A\mathbf{x}$$
. Simplify to the extent possible.  

$$A = \begin{bmatrix} 1 & 0 & -3 \\ -2 & -1 & 4 \end{bmatrix} \quad \mathbf{x} = \begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix} \qquad \overset{n \sim 2}{\overset{n \sim 2}{\underset{-1}{1}} \qquad \overset{n \sim 2}{\underset$$

Find the product *A***x**. Simplify to the extent possible.

$$A = \begin{bmatrix} 2 & 4 \\ -1 & 1 \\ 0 & 3 \end{bmatrix} \quad \mathbf{x} = \begin{bmatrix} -3 \\ 2 \end{bmatrix} \qquad \begin{array}{c} \mathbf{m} = 3 \\ \vec{x} \text{ in } \mathbb{R}^2 \\ \vec{x} \text{ in } \mathbb{R}^2 \\ A \vec{x} = x_1 \vec{a}_1 + x_2 \vec{a}_2 \\ = -3 \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 4 \\ -3 \\ 2 \end{bmatrix} = \begin{bmatrix} -3 \\ -3 \end{bmatrix} + \begin{bmatrix} 2 \\ -1 \\ -3 \end{bmatrix} + \begin{bmatrix} 2 \\ -1 \\ -3 \end{bmatrix} + \begin{bmatrix} 2 \\ -1 \\ -3 \end{bmatrix} = \begin{bmatrix} 2 \\ -3 \\ -3 \end{bmatrix} = \begin{bmatrix} 2 \\$$

▲□▶▲圖▶★厘▶★厘▶ 厘 のQC

0-2

Write the linear system as a vector equation and then as a matrix equation of the form  $A\mathbf{x} = \mathbf{b}$ .



#### Theorem

If *A* is the  $m \times n$  matrix whose columns are the vectors  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$ , and **b** is in  $\mathbb{R}^m$ , then the matrix equation

$$A\mathbf{x} = \mathbf{b}$$

has the same solution set as the vector equation

$$x_1a_1 + x_2a_2 + \cdots + x_na_n = b$$

which, in turn, has the same solution set as the linear system of equations whose augmented matrix is

$$[\mathbf{a}_1 \ \mathbf{a}_2 \ \cdots \ \mathbf{a}_n \ \mathbf{b}].$$



The equation  $A\mathbf{x} = \mathbf{b}$  has a solution if and only if **b** is a linear combination of the columns of *A*.

In other words, the corresponding linear system is consistent if and only if **b** is in Span{ $\mathbf{a}_1, \mathbf{a}_2, \ldots, \mathbf{a}_n$  }.

Characterize the set of all vectors  $\mathbf{b} = (b_1, b_2, b_3)$  such that  $A\mathbf{x} = \mathbf{b}$  has a solution where

$$A = \left[ \begin{array}{rrrr} 1 & 3 & 4 \\ -4 & 2 & -6 \\ -3 & -2 & -7 \end{array} \right].$$

be can determine the b's by using equivalence to the system with augmented matrix [Ab]. Will use row reduction. Find the met of

$$\begin{bmatrix} 1 & 3 & 4 & b_1 \\ -4 & 2 & -6 & b_2 \\ -3 & -2 & -7 & b_3 \end{bmatrix}$$

$$4R_1 + R_2 \rightarrow R_2$$
$$3R_1 + R_3 \rightarrow R_3$$

(4日)(4日)(4日)(4日)(日)(900)

◆□▶ ◆圖▶ ◆目▶ ◆目▶ 目 - のへで

9/35

The system is consistent only if the 4th column is not a pivot column. The system is consisten if  $2b_{3}+6b_{1}-(b_{2}+4b_{1})=0$  $3b_1 - b_2 + 2b_3 = 0$ The system is consistent provided the entries of I satisfy this condition. We can state  $ab_{1} = b_{2} - 2b_{3}$ this as  $b_1 = \frac{1}{2}b_2 - b_3$ where by and by an free

The vectors look like  

$$\vec{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} = \begin{bmatrix} \frac{1}{2}b_2 & -b_3 \\ b_2 \\ b_3 \end{bmatrix} = \begin{bmatrix} \frac{1}{2}b_2 \\ b_3 \\ b_3 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} +$$

## Theorem (first in a string of equivalency theorems)

Let *A* be an  $m \times n$  matrix. Then the following are logically equivalent (i.e. they are either all true or are all false).

(a) For each **b** in  $\mathbb{R}^m$ , the equation  $A\mathbf{x} = \mathbf{b}$  has a solution.

(b) Each **b** in  $\mathbb{R}^m$  is a linear combination of the columns of *A*.

(c) The columns of A span  $\mathbb{R}^m$ .

(d) A has a pivot position in every row.

(Note that statement (d) is about the *coefficient* matrix A, not about an augmented matrix  $\begin{bmatrix} A & \mathbf{b} \end{bmatrix}$ .)

# Computing Ax

We can use a *row-vector* dot product rule. The  $i^{th}$  entry is  $A\mathbf{x}$  is the sum of products of corresponding entries from row i of A with those of  $\mathbf{x}$ . For example

$$\begin{bmatrix} 1 & 0 & -3 \\ -2 & -1 & 4 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix} : \begin{bmatrix} 1 \cdot 2 + 0 \cdot | + (\cdot 3) \cdot (-1) \\ -2 \cdot 2 + (-1) \cdot | + 4 \cdot (-1) \end{bmatrix} : \begin{bmatrix} 5 \\ -9 \end{bmatrix}$$

< 日 > < 同 > < 回 > < 回 > < 回 > <

 $\begin{bmatrix} 2 & 4 \\ -1 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} -3 \\ 2 \end{bmatrix} = \begin{bmatrix} 2(-3) + 4(2) \\ -1(-3) + 1(2) \\ 0(-3) + 3(2) \end{bmatrix} = \begin{bmatrix} 2 \\ 5 \\ 6 \end{bmatrix}$ 3x 2 R<sup>2</sup>

 $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1x_1 + 0x_2 + 0x_3 \\ 0x_1 + 1x_2 + 0x_3 \\ 0x_1 + 0x_2 + 1x_3 \end{bmatrix} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$   $3 \times 3 \quad \mathbb{R}^3$ 

14/35

## **Identity Matrix**

We'll call an  $n \times n$  matrix with 1's on the diagonal and 0's everywhere else—i.e. one that looks like

$$\begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}$$

the  $n \times n$  **identity** matrix and denote it by  $I_n$ . (We'll drop the subscript if it's obvious from the context.)

This matrix has the property that for each  $\mathbf{x}$  in  $\mathbb{R}^n$ 

$$l_n \mathbf{x} = \mathbf{x}.$$

< 日 > < 同 > < 回 > < 回 > < 回 > <

#### Theorem: Properties of the Matrix Product

If *A* is an  $m \times n$  matrix, **u** and **v** are vectors in  $\mathbb{R}^n$ , and *c* is any scalar, then

(a)  $A(\mathbf{u} + \mathbf{v}) = A\mathbf{u} + A\mathbf{v}$ , and

(b)  $A(c\mathbf{u}) = cA\mathbf{u}$ .